arXiv:hep-th/0004007v1 3 Apr 2000

FIELD-DEPENDENT BRS TRANFORMATIONS AND CORRECT PRESCRIPTION FOR $\frac{1}{(\eta\cdot k)^p}$-TYPE SINGULARITIES IN AXIAL GAUGES

A. Misra
Institute of Physics, Sachivalaya Marg, Bhubaneswar 751005, India



The axial-gauge boson propagator contains $1/(\eta\cdot k)^p$-type singularities. These singualarities have generally been treated by inventing prescriptions for them. We propose an alternative procedere for treating these singularities in the path-integral formalism using the known way of treating the $1/k^{2n}$-type singularities in Lorentz-type gauges. For this purpose we use a finite field-dependent BRS transformation that inerpolates between the Lorentz and axial-type gauges. We arrive at the ϵ-dependent tree propagator in axial-type gauges.

Calculations in nonabelaian gauge theories require a choice of gauge. There are many families of gauges that have been used in practical calculations, e.g., Lorentz-type gauges, axial gauges. *It becomes an important question as to how the calculations in various (families of) gauge choices are related to each other.* Gauge-independence in a limited framework, has been proven in early days. Such proofs utililize the *infinitesimal* gauge transformations reponsible for gauge-parameter change. Ways of connecting Green's functions in family of gauges (and establishing expliticly gauge-independence) has not been done until recently. Discrepancies have been reported in anomalous dimension calculations in the Lorentz-type and axial-type gauges. Thus, it becomes important to obtain a procedure to connect the Green functions in different families of gauges.

Unlike gauge transformations, "infinitesimal" and "finite" BRS transformations have the same form. This makes them suitable for our purpose. The FP effective actions in the Lorentz and Axial gauges are invariant under the BRS transformations:

$$\delta\phi_i(x) = \delta_{\rm BRS}\phi_i(x)\delta\Lambda \tag{1}$$

where $\delta_{\rm BRS}\phi_i(x)$ is given in [1)], and the gauge and (anti)ghost fields are generically denoted by ϕ_i.

As observed by Joglekar and Mandal [1], $\delta\Lambda$ *need not be infinitesimal nor need it be field-independent as long as it does not depend on x explicitly for (1) to be a symmetry of FPEA* In fact, the following finite field-dependent BRS (FFBRS) transformations were introduced:

$$\phi_i'(x) = \phi_i(x) + \delta_{\rm BRS}\phi_i(x)\Theta[\phi], \tag{2}$$

where $\Theta[\phi]$ is an x-independent functional of $A,\ c,\ \bar{c}$ and these were also the symmetry of the FPEA.

The FPEA is invariant under (1), but the functional measure is not invariant under the (nonlocal) transformations (1). The Jacobian for the FFBRS transformations can be expressed (in special cases effectively as $exp(iS_1)$ and this S_1 explains the difference between the two effective actions [1), 2)].

Such FFBRS transformations were constructed by integration of an infinitesimal field-dependent BRS (IFBRS) transformation:

$$\frac{d\phi_i(x,\kappa)}{d\kappa} = \delta_{\rm BRS}[\phi(x,\kappa)]\Theta'[\phi(x,\kappa)]. \tag{3}$$

The integration of (3) from $\kappa = 0$ to 1, leads to the FFBRS transformation of (2) with $\phi(\kappa = 1) \equiv \phi'$ and $\phi(\kappa = 0) = \phi$. The relationship between Θ and Θ' is given in ref 1.

The result for the FFBRS Transformation for Correlating Green's functions in Lorentz-type Gauges to Axial-type Gaugees (that differs from the FFBRS of refs. 1 and 2) reads [3)]:

$$\begin{aligned}\langle O[\phi]\rangle_A &\equiv \int \mathcal{D}\phi' O[\phi'] e^{iS^A_{\rm eff}[\phi']} \\ &= \int \mathcal{D}\phi \mathcal{O}[\phi] e^{iS^L_{\rm eff}[\phi]} + \int \mathcal{D}\phi \sum_i \delta\phi_i[\phi]\frac{\delta\mathcal{O}}{\delta\phi_i} e^{iS^L_{\rm eff}},\end{aligned} \tag{4}$$

where the summation over i runs over fields $A, c, \bar{c}$ and

$$\begin{aligned}\phi' &= \phi + \Big(\tilde{\delta}_1[\phi]\Theta_1[\phi] + \tilde{\delta}_2[\phi]\Theta_2[\phi]\Big)\Theta'[\phi] \\ &\equiv \phi + \delta\phi[\phi]\end{aligned} \tag{5}$$

is an FFBRS with

$$\Theta_{1,2}[\phi] \equiv \int_0^1 d\kappa(1,\kappa) exp\Big(\kappa f_1[\phi] + \frac{\kappa^2}{2} f_2[\phi]\Big); \tag{6}$$

$$f[\tilde{\phi},\kappa] \equiv f_1[\tilde{\phi}] + \kappa f_2[\tilde{\phi}]; \tag{7}$$

$$\begin{aligned}f_1[\phi] &\equiv i\int d^4x \Big[\frac{\partial\cdot A^\alpha}{\lambda}(\partial\cdot A^\alpha - \eta\cdot A^\alpha) + \bar{c}(\partial\cdot \mathrm{D} - \eta\cdot \mathrm{D})c\Big] \\ f_2[\phi] &\equiv -\frac{i}{\lambda}\int d^4x(\partial\cdot A^\alpha - \eta\cdot A^\alpha)^2,\end{aligned} \tag{8}$$

and

$$\Theta' \equiv i\int d^4x\ \bar{c}^\alpha(\partial\cdot A^\alpha - \eta\cdot A^\alpha). \tag{9}$$

An alternate and more effective expression can be given:

$$\langle\mathcal{O}\rangle_A = \langle\mathcal{O}\rangle_L + \int_0^1 d\kappa \int D\phi \sum_i \Big(\tilde{\delta}_{1,i}[\phi] + \kappa\tilde{\delta}_{2,i}[\phi]\Big)\Theta'[\phi]\frac{\delta\mathcal{O}}{\delta\phi_i} e^{iS^M_{\rm eff}}, \tag{10}$$

where $\tilde{\delta}_{1,i}$ and $\tilde{\delta}_{2,i}$ are defined in via

$$\tilde{\delta}_{\rm BRS}\phi_i \equiv \Big(\tilde{\delta}_{1,i} + \kappa\tilde{\delta}_{2,i}\Big)\delta\Lambda \tag{11}$$

where $\tilde{\delta}_{\rm BRS}\phi_i$ are the BRS variations for the mixed gauge function $[\partial\cdot A(1-\kappa)+\kappa\eta\cdot A]$.

The basic idea is to use (10) to relate the axial and Lorentz gauge propagators. The only shortcoming of the above relation is that it does not include the $i[-\epsilon A^2/2+\epsilon\bar{c}c]$ terms in the Lorentz gauge effective action. The modification of (10) is:

$$\begin{aligned}&\langle O\rangle_A = \langle O\rangle_L \\ &+\int_0^1 d\kappa\int \mathcal{D}\phi e^{i[S^M_{\rm eff}[\phi,\kappa]-i\epsilon(A^2/2-\bar{c}c)]}\Big[\Big(\delta_1[\phi]+\kappa\delta_2[\phi]\Big)\Theta'[\phi]\frac{\delta O}{\delta\phi}\Big].\end{aligned} \tag{12}$$

Thus, in *this form*, the only effect on the second term is to modify $S^M_{\rm eff}$ by $-i\epsilon\int(A^2/2-\bar{c}c)d^4x$ inside κ-integration.

We now employ the result (12) for the propagators. We set $O[\phi]=A^\alpha_\mu(x)A^\beta_\nu(y)$. The equation (12) then reads:

$$\begin{aligned}&iG^{0A\ \alpha\beta}_{\mu\nu}(x-y) = iG^{0L\ \alpha\beta}_{\mu\nu}(x-y) + i\int_0^1 d\kappa\int\mathcal{D}\phi e^{i[S^M_{\rm eff}[\phi,\kappa]-i\epsilon\int(A^2/2-\bar{c}c)d^4x]} \\ &\times\Big((\mathrm{D}_\mu c)^\alpha(x)A^\beta_\nu(y)+A^\alpha_\mu(x)(\mathrm{D}_\nu c)^\beta(y)\Big)\int d^4z\bar{c}^\gamma(z)(\partial\cdot A^\gamma-\eta\cdot A^\gamma)(z).\end{aligned}$$

This leads to, for zero loop case [4)],

$$\tilde{G}^{0A}_{\mu\nu} = \tilde{G}^{0L}_{\mu\nu} + \Big[\Big(k_\mu k_\nu\Sigma_1+\eta_\mu k_\nu\Sigma_2\Big)ln\Sigma_3 + (k\to -k;\mu\leftrightarrow\nu)\Big] \tag{13}$$

where

$$\begin{aligned}\Sigma_1 &\equiv \frac{-(k^2-i\eta\cdot k)\Big(\frac{\eta\cdot k+i\eta^2}{k^2-i\eta\cdot k}+i\lambda-\frac{(1-\lambda)\eta\cdot k}{k^2+i\epsilon}\Big)}{\epsilon\Sigma} \\ \Sigma_2 &\equiv \frac{-(k^2-i\eta\cdot k)\Big(-\Big[\frac{k^2+i\eta\cdot k}{k^2-i\eta\cdot k}\Big]+1-\frac{i\epsilon(1-\lambda)}{k^2+i\epsilon}\Big)}{\epsilon\Sigma} \\ \Sigma_3 &\equiv \frac{-i(\eta\cdot k+\epsilon)(k^2+i\epsilon\lambda)}{(k^2+i\epsilon)\Big(-i\epsilon\lambda-\sqrt{k^4-(k^2+i\epsilon\lambda)\Big[k^2+\frac{(\eta\cdot k)^2+i\epsilon\eta^2}{k^2+i\epsilon}\Big]}\Big)}, \\ &\text{and} \\ \Sigma &\equiv \Big[(1-\lambda)[(\eta\cdot k)^2+2ik^2\eta\cdot k]+i\epsilon k^2(1-2\lambda)+\lambda(k^2+i\epsilon)^2+\eta^2(k^2+i\epsilon)\Big].\end{aligned} \tag{14}$$

The k^0 integration over this propagator can be replaced by a k^0-integration over (most of) the real axis combined over semicircle in the LHP of radius $>>\sqrt{\epsilon}$ (where the complication due to presence of ϵ can be dropped and the usual simple form can be used) and an additional effective term of much simpler form that rounds up effectively the complex structure near $\eta\cdot k=0$ [5)]. For $\eta^2\neq 0$, and $k^2\neq 0$, the latter reads

$$\delta\Big(k^0-\frac{1}{2}\sqrt{\frac{\epsilon\eta^2}{i}}-\vec{\eta}\cdot\vec{k}\Big)\Big[k_\mu k_\nu D_1+\eta_\mu k_\nu D_2+\eta_\nu k_\mu D_3\Big]+\mu\leftrightarrow\nu; k\to -k \tag{15}$$

$$D_1 \equiv -\frac{\pi\eta^2}{\mathcal{K}_1} i\sqrt{\frac{i\eta^2}{\epsilon}} + \frac{i\pi(\eta^2)^2\mathcal{K}_2}{2\mathcal{K}_1^2};$$
$$D_2 \equiv \frac{i\pi\eta^2}{2\mathcal{K}_1};\ D_3 \equiv -\frac{i\pi\eta^2}{2\mathcal{K}_1}, \tag{16}$$

where

$$\mathcal{K}_1 \equiv \Big((\vec{\eta}\cdot\vec{k})^2 - \vec{k}^2\Big)(\eta^2 + i\epsilon);$$
$$\mathcal{K}_2 \equiv 2i\Big((\vec{\eta}\cdot\vec{k})^2 - \vec{k}^2\Big) + 2\vec{\eta}\cdot\vec{k}(\eta^2 + i\epsilon). \tag{17}$$

Note *that if we define the LCG as the* $\eta^2 \to 0$ *limit, then this additional term (15) vanishes.* Thus, we obtain a simple result of the LCG.

For $|\eta\cdot k| >> \epsilon$, (13) gives the expected result:

$$\tilde{G}^{0A}_{\mu\nu} - \tilde{G}^{0L}_{\mu\nu} = -\frac{1}{k^2}k_\mu k_\nu\Big(\frac{(\lambda k^2+\eta^2)}{(\eta\cdot k)^2} + \frac{(1-\lambda)}{k^2}\Big) + \frac{k_{[\mu}\eta_{\nu]_+}}{k^2\eta\cdot k}. \tag{18}$$

As the FFBRS transformations discussed also preserve the vacuum expectation values of gauge-invariant observables, it follows that our treatment is such that by its very construction, the Wilson loop $W[L]$ has the same value in the Lorentz and axial-type gauges to *all* orders. We have further given the proof of this statement to $O(g^4)$ using the earlier work by Cheng and Tsai. Our proof holds for any arbitrary loop for $\eta^2 < 0$ and for a subclass of loops for $\eta^2 \geq 0$ [6].

We also note that the $O(g^2)$ thermal Wilson loop W_R [6], depends only on $D_{00}(k^0 = 0, \vec{k})$. We find that for the propagator in (13), $D_{00}(k^0 = 0, \vec{k}) = \frac{g_{00}}{k^2+i\epsilon}$ which is the same as $D_{00}(k^0 = 0, \vec{k})$ for Lorentz gauges and as such W_R has the same value as in Lorentz-type gauges.

In conclusion, we addressed the problem of relating arbitrary Green's functions in two sets of uncorrelated gauges, e.g.. the axial and the Lorentz-type gauges (the example considered). We showed that this involved an FFBRS, obtained by intregration of an IFBRS. We found that the final result could be put in a neat form (4) or (10). Using our result, we have *derived* the correct prescription for Axial gauge propagator. Even though, the propagator in axial gauge, naively calculated, has spurious singularities. the correct treatment of these singularities is obtained by relating this propagator to the corresponding Lorentz gauge treatment. This was done by using the FFBRS discussed in this talk. The propagator of (13) gives, however complex, the actual correct treatment of these singularities. While for $|\eta\cdot k| >> \epsilon$, it gives the usual propagator, the actual analytic nature of the propagator, in the vicinity of the origin is much more complicated than indicated by various prescriptions suggested earlier.

Acknowledgement

I wish to thank my collaborator Prof.S.D.Joglekar (IIT Kanpur, India) for introducing me to FFBRS transformations and sharing his insights with me. I also wish to thank the Moriond committee for inviting me to the conference.

References

1. S.D.Joglekar and B.P.Mandal, Phys Rev D **51**, 1919 (1995).
2. R.S.Bandhu and S.D.Joglekar J.Phys.A-Math and General **31**, 4217 (1998).
3. S.D.Joglekar and A.Misra, J. of Math. Phys **41**, 1755 (2000).
4. S.D.Joglekar and A.Misra, Mod. Phys. Lett A **14**, No. 30, 2083 (1999).
5. S.D.Joglekar and A.Misra, to appear in Int. J. of Mod. Phys A (2000).
6. S.D.Joglekar and A.Misra, to appear in Mod. Phys. Lett A (2000).